




# Cross Section Measurements

## Ingredients to a Cross Section

*Mostly Deep Inelastic Scattering*





# Prerequisites and Reminders ...

Natural Units

Four-Vector Kinematics

Lorentz Transformation

Lorentz Boost

Lorentz Invariance

Rapidity etc.

Invariant Mass

CMS-Energy

Particle Decays

Cross Section

Matrix Element

Phase Space

Feynman Diagrams

Mandelstam Variables

Parton Distributions

Bjorken-x

...

$$\hbar = 1, c = 1$$

$$\hbar c = 197.3 \text{ MeV fm}$$

$$(\hbar c)^2 = 0.3894 \text{ GeV}^2 \text{ mb}$$

$$p = (E, \vec{p})$$

$$p^2 = E^2 - \vec{p}^2 = m^2$$

$$\beta = p/E, \gamma = E/m$$

$$p_1 \cdot p_2 = E_1 E_2 - \vec{p}_1 \cdot \vec{p}_2$$

4-vector scalar product

Lorentz invariant

→ All quantities like cross sections etc.  
should be in terms of scalar products of 4-vectors ...





# Prerequisites and Reminders ...

Natural Units  
Four-Vector Kinematics  
Lorentz Transformation  
Lorentz Boost  
Lorentz Invariance  
Rapidity etc.  
Invariant Mass  
CMS-Energy  
  
Particle Decays  
Cross Section  
Matrix Element  
Phase Space  
Feynman Diagrams  
Mandelstam Variables  
  
Parton Distributions  
Bjorken-x  
...

$$p = (E, \vec{p})$$

Particle momentum as seen  
in laboratory frame ...

$$p^* = (E^*, \vec{p}^*)$$

Particle momentum as viewed from a  
frame moving with velocity  $\beta_f$  ...

Lorentz Transformation:

$$E^* = \gamma_f \cdot E - \gamma_f \beta_f \cdot p_{\parallel}$$

$$p_{\parallel}^* = \gamma_f \cdot p_{\parallel} - \gamma_f \beta_f \cdot E$$


$$p_T^* = p_T$$

$$\text{with } \gamma_f = (1 - \beta_f^2)^{-\frac{1}{2}}$$



# Prerequisites and Reminders ...

$p_T$  is conserved,  $p_l$  of colliding objects is unknown  
 $\rightarrow$  constraint only in transverse plane




Schematic  
proton-proton scattering

Relevant kinematic variables:

- Transverse momentum:  $p_T$
- Rapidity:  $y = \frac{1}{2} \ln (E-p_z)/(E+p_z)$
- Pseudorapidity:  $\eta = -\ln \tan \frac{1}{2}\theta$
- Azimuthal angle:  $\phi$

Pseudorapidity




Physics at Hadron Colliders

Natural Units  
 Four-Vector Kinematics  
 Lorentz Transformation  
 Lorentz Boost  
 Lorentz Invariance  
 Rapidity etc.  
 Invariant Mass  
 CMS-Energy  
 Particle Decays  
 Cross Section  
 Matrix Element  
 Phase Space  
 Feynman Diagrams  
 Mandelstam Variables  
 Parton Distributions  
 Bjorken-x  
 ...



# Prerequisites and Reminders ...

Natural Units  
 Four-Vector Kinematics  
 Lorentz Transformation  
 Lorentz Boost  
 Lorentz Invariance  
 Rapidity etc.  
 Invariant Mass  
 CMS-Energy  
  
 Particle Decays  
 Cross Section  
 Matrix Element  
 Phase Space  
 Feynman Diagrams  
 Mandelstam Variables  
  
 Parton Distributions  
 Bjorken-x  
 ...



$$\begin{aligned}
 M^2 &= (p_1 + p_2)^2 \\
 &= (E_1 + E_2)^2 - (\vec{p}_1 + \vec{p}_2)^2 \\
 &= m_1^2 + m_2^2 + 2E_1 E_2 (1 - \vec{\beta}_1 \vec{\beta}_2)
 \end{aligned}$$

Center-of-mass Energy:

$$E_{\text{cm}} = \left[ (E_1 + E_2)^2 - (\vec{p}_1 + \vec{p}_2)^2 \right]^{\frac{1}{2}}$$

Particle 2 at rest:

$$E_{\text{cm}} = \left[ m_1^2 + m_2^2 + 2E_1 m_2 \right]^{\frac{1}{2}}$$

Particle Collider:


$$[E_1 = E_2; \vec{p}_1 = -\vec{p}_2; m_1 = m_2 \approx 0]$$

$$E_{\text{cm}} = 2E$$





# Prerequisites and Reminders



Natural Units  
 Four-Vector Kinematics  
 Lorentz Transformation  
 Lorentz Boost  
 Lorentz Invariance  
 Rapidity etc.  
 Invariant Mass  
 CMS-Energy

Cross Section  
 Particle Decays  
 Matrix Element  
 Phase Space

Feynman Diagrams  
 Mandelstam Variables

Parton Distributions  
 Bjorken-x

...

Differential  
Cross Section:

$$d\sigma = \frac{(2\pi)^4 |\mathcal{M}|^2}{4\sqrt{(p_1 \cdot p_2)^2 - m_1^2 m_2^2}} \times d\Phi_n(p_1 + p_2; p_3, \dots, p_{n+2})$$

Physics

Kinematics

Matrix element

n-body phase space


$$d\Phi_n = \dots$$

$$\dots = \delta^4(P - \sum_{i=1}^n p_i) \prod_{i=1}^n \frac{d^3 p_i}{(2\pi)^3 2E_i}$$

with  $P = p_1 + p_2$

Partial  
Decay Rate:


$$d\Gamma = \frac{(2\pi)^4}{2M} |\mathcal{M}|^2 \times d\Phi_n(P; p_1, \dots, p_n)$$





# Prerequisites and Reminders ...

Natural Units  
 Four-Vector Kinematics  
 Lorentz Transformation  
 Lorentz Boost  
 Lorentz Invariance  
 Rapidity etc.  
 Invariant Mass  
 CMS-Energy  
 Particle Decays  
 Cross Section  
 Matrix Element  
 Phase Space  
 Feynman Diagrams  
 Mandelstam Variables  
 Parton Distributions  
 Bjorken-x  
 ...






Mandelstam variables:

$$\begin{aligned}
 s &= (p_1 + p_2)^2 = (p_3 + p_4)^2 \\
 t &= (p_1 - p_3)^2 = (p_2 - p_4)^2 \\
 u &= (p_1 - p_4)^2 = (p_2 - p_3)^2
 \end{aligned}$$

Relativistic limit ( $m_i$  is the mass of particle  $i$ ):

$$\begin{aligned}
 s &\approx 2p_1 \cdot p_2 \approx 2p_3 \cdot p_4 \\
 t &\approx -2p_1 \cdot p_3 \approx -2p_2 \cdot p_4 \\
 u &\approx -2p_1 \cdot p_4 \approx -2p_3 \cdot p_2
 \end{aligned}$$

$$s + t + u = m_1^2 + m_2^2 + m_3^2 + m_4^2$$





# Prerequisites and Reminders

Natural Units  
Four-Vector Kinematics  
Lorentz Transformation  
Lorentz Boost  
Lorentz Invariance  
Rapidity etc.  
Invariant Mass  
CMS-Energy

Particle Decays  
Cross Section  
Matrix Element  
Phase Space  
Feynman Diagrams  
Mandelstam Variables

Parton Distributions  
Bjorken-x

...

Proton-Proton  
Cross Section:


$$\sigma = \sum_{ij} \int dx_1 dx_2 f_i(x_1, Q^2) f_j(x_2, Q^2) \hat{\sigma}(Q^2)$$

Parton content:  
 $f(x, Q^2) = q(x, Q^2)$  or  $g(x, Q^2)$

$x_{1,2}$  : Bjorken-x  
fractional momentum of parton  
involve in hard process

$Q^2$  : scale; spacial resolution  
Invariant Parton Parton mass


$f$  : Parton Distribution function  
measured e.g. at HERA ...








# Proton-Proton Scattering @ LHC





# Feynman Diagrams



Fermi's Golden Rule

$$W_{\text{fi}} = 2\pi |M_{\text{fi}}|^2 \cdot \frac{dN}{dE_{\text{f}}}$$

Transition  
probability

Matrix element

Phase space

4-vector current










$$M_{\text{fi}} = -i \int j_{\mu}^{(1)} \cdot \left( \frac{1}{q^2} \right) \cdot j_{\mu}^{(2)} d^4x$$

Propagator

$$\begin{aligned} \sigma &\sim |M_{\text{fi}}|^2 \\ &\sim g^4 \cdot \left( \frac{1}{q^4} \right) \end{aligned}$$



# Some Hard Processes with quarks and gluons ...






# QCD Matrix Elements

Hard scattering with quarks and gluons!


$g_s$  = QCD coupling strength



Subprocess	$ \mathcal{M} ^2/g_s^4$	$ \mathcal{M}(90^\circ) ^2/g_s^4$
$qq' \rightarrow qq'$ $q\bar{q}' \rightarrow q\bar{q}'$	$\frac{4}{9} \frac{\hat{s}^2 + \hat{u}^2}{\hat{t}^2}$	2.2
$qq \rightarrow qq$	$\frac{4}{9} \left( \frac{\hat{s}^2 + \hat{u}^2}{\hat{t}^2} + \frac{\hat{s}^2 + \hat{t}^2}{\hat{u}^2} \right) - \frac{8}{27} \frac{\hat{s}^2}{\hat{u}\hat{t}}$	3.3
$q\bar{q} \rightarrow q'\bar{q}'$	$\frac{4}{9} \frac{\hat{t}^2 + \hat{u}^2}{\hat{s}^2}$	0.2
$q\bar{q} \rightarrow q\bar{q}$	$\frac{4}{9} \left( \frac{\hat{s}^2 + \hat{u}^2}{\hat{t}^2} + \frac{\hat{t}^2 + \hat{u}^2}{\hat{s}^2} \right) - \frac{8}{27} \frac{\hat{u}^2}{\hat{s}\hat{t}}$	2.6
$q\bar{q} \rightarrow gg$	$\frac{32}{27} \frac{\hat{u}^2 + \hat{t}^2}{\hat{u}\hat{t}} - \frac{8}{3} \frac{\hat{u}^2 + \hat{t}^2}{\hat{s}^2}$	1.0
$gg \rightarrow q\bar{q}$	$\frac{1}{6} \frac{\hat{u}^2 + \hat{t}^2}{\hat{u}\hat{t}} - \frac{3}{8} \frac{\hat{u}^2 + \hat{t}^2}{\hat{s}^2}$	0.1
$qg \rightarrow qg$	$\frac{\hat{s}^2 + \hat{u}^2}{\hat{t}^2} - \frac{4}{9} \frac{\hat{s}^2 + \hat{u}^2}{\hat{u}\hat{s}}$	6.1
$gg \rightarrow gg$	$\frac{9}{4} \left( \frac{\hat{s}^2 + \hat{u}^2}{\hat{t}^2} + \frac{\hat{s}^2 + \hat{t}^2}{\hat{u}^2} + \frac{\hat{u}^2 + \hat{t}^2}{\hat{s}^2} + 3 \right)$	30.4



# Proton-Proton Scattering @ LHC






# Electron-Proton Scattering

PDFs are needed to compute cross-section


→ How to measure PDFs? Unfolding 2 PDFs is ~difficult → **replace p with e!**








# Electron-Proton Scattering @ HERA





# HERA




HERA was an Hadron-Electron Ring Accelerator in Desy, Hamburg-DE. It began operating in 1992 around 15 to 30 m underground and has a circumference of 6.3 km. . At HERA, electrons or positrons were collided with protons at a cms energy of 318 GeV. HERA was closed down on 30 June 2007. The HERA tunnel is located under the DESY site and the nearby Volkspark Leptons and protons were stored in two independent storage rings on top of each other inside this tunnel. There are four interaction regions, which were used by the experiments H1, ZEUS, HERMES and Hera-B. Leptons (electrons or positrons) were pre-accelerated to 450 MeV in the linear accelerator *LINAC-II*. From there they were injected into the storage ring *DESY-II* and accelerated further to 7.5 GeV before their transfer into *PETRA*, where they were accelerated to 14 GeV. Finally they were injected into their storage ring in the HERA tunnel and reached a final energy of 27.5 GeV. This storage ring was equipped with warm (non-superconducting) magnets keeping the leptons on their circular track by a magnetic field of 0.17 Tesla. Protons were obtained from originally negatively charged hydrogen ions and pre-accelerated to 50 MeV in a linear accelerator. They were then injected into the proton synchrotron *DESY-III* and accelerated further to 7 GeV. Then they were transferred to *PETRA* where they were accelerated to 40 GeV. Finally, they were injected into their storage ring in the HERA tunnel and reached their final energy of 920 GeV. The proton storage ring used superconducting magnets to keep the protons on track. The characteristic build-up time expected for the HERA accelerator was approximately 40 minutes.





# HERA




HERA: pe Accelerator in Desy, Hamburg-DE:

- It began operating in 1992. and was closed down in 2007.
- circumference of 6.3 km
- Leptons and protons were stored in two independent storage rings on top of each other inside this tunnel.
- warm (non-superconducting) with a magnetic field of 0.17 Tesla for leptons  
superconducting magnets for protons
- electrons (or positrons): 450 MeV (*LINAC-II*) → 7.5 GeV (*DESY-II*) → 14 GeV (*PETRA*)  
→ 27.5 GeV (HERA)
- Protons : 50 MeV (Linac) → to 7 GeV (*DESY-III*) → 40 GeV (*PETRA*) → 920 GeV (HERA).
- final cms energy: 318 GeV
- four interaction regions, → experiments H1, ZEUS, HERMES and Hera-B.



# Hera Accelerator at Desy








# Electron-Proton Scattering @ HERA


Asymmetric beams  $\rightarrow$  unbalanced event, mostly in the p direction  
 $\rightarrow$  asymmetric detector








# Electron-Proton Scattering @ HERA




# Structure Function $F_2$




$$\frac{d\sigma(eq)}{dq^2} = \frac{4\pi\alpha^2}{q^4} e_q^2$$

Rutherford scattering  
on pointlike target



$$\frac{d\sigma(ep)}{dq^2} = \frac{4\pi\alpha^2}{q^4} [2e_u^2 + e_d^2] = \frac{4\pi\alpha^2}{q^4}$$

With  
quark-quark  
interactions




$$\begin{aligned} \frac{d\sigma(ep)}{dx dq^2} &= \frac{4\pi\alpha^2}{q^4} [e_u^2 u(x) + e_d^2 d(x) + \dots] \\ &= \frac{4\pi\alpha^2}{q^4} \frac{F_2(x)}{x} \end{aligned}$$


QPM: Structure Functions  $F_2$  independent of  $Q^2$




Proton Three  
valence quarks



Proton Three bound  
valence quarks






Proton Bound valence  
quarks + gluon radiation



[see e.g. Halzen/Martin]


small x

$F_2(x)$





# Invariant Quantities in DIS



Invariant quantities:

$\nu = \frac{q \cdot P}{M} = E - E'$  is the lepton's energy loss in the nucleon rest frame (in earlier literature sometimes  $\nu = q \cdot P$ ). Here,  $E$  and  $E'$  are the initial and final lepton energies in the nucleon rest frame.

$Q^2 = -q^2 = 2(EE' - \vec{k} \cdot \vec{k}') - m_\ell^2 - m_{\ell'}^2$ , where  $m_\ell(m_{\ell'})$  is the initial (final) lepton mass. If  $EE' \sin^2(\theta/2) \gg m_\ell^2, m_{\ell'}^2$ , then  $Q^2 \approx 4EE' \sin^2(\theta/2)$ , where  $\theta$  is the lepton's scattering angle with respect to the lepton beam direction.

$x = \frac{Q^2}{2M\nu}$  where, in the parton model,  $x$  is the fraction of the nucleon's momentum carried by the struck quark.

$y = \frac{q \cdot P}{k \cdot P} = \frac{\nu}{E}$  is the fraction of the lepton's energy lost in the nucleon rest frame.

$W^2 = (P + q)^2 = M^2 + 2M\nu - Q^2$  is the mass squared of the system  $X$  recoiling against the scattered lepton.

$s = (k + P)^2 = \frac{Q^2}{xy} + M^2 + m_\ell^2$  is the center-of-mass energy squared of the lepton-nucleon system.



# Kinematics of DIS - 1

Particle Data Group  
– PDG CERN

## 8. Deep inelastic scattering.

In lepton-hadron scattering at sufficiently high energies one finds a large number of hadrons in the final state: this is *deep inelastic scattering* (DIS). The multiplicity of the hadronic system varies event by event. The reaction equation for electron-proton DIS is written as

$$e^- + p \rightarrow e^- + X \quad (84)$$

where  $X$  stands for the hadronic system with an arbitrary number of particles. A generic diagram depicting the DIS process is shown in Fig. 2.




Figure 2: Generic diagram of deep inelastic scattering.

To describe the DIS reaction kinematics we denote the 4-momentum of the incoming electron by  $k = (E, 0, 0, k)$ , that of the target proton by  $P$  and those of the scattered electron and of the hadronic system by  $k'$  and  $P'$ , respectively. The exchanged virtual photon  $\gamma^*$  has 4-momentum  $q = k - k'$ . 4-momentum conservation demands

$$k + P = k' + P' \quad (85)$$

and we have the mass-shell conditions  $k^2 = k'^2 = m_e^2$  and  $P^2 = m_p^2$ . Since energies characteristic of DIS are at least of several GeV, the electron mass can be safely set equal to zero. Then we get for the square of the 4-momentum transfer  $q^2 = (k - k')^2 = -2EE'(1 - \cos \theta)$ , and we see that  $q^2 \leq 0$ , *i.e.* the exchanged photon is space-like.





# Kinematics of DIS - 2

The invariant  $W^2 = P'^2$  is variable because of the variable multiplicity of particles in the hadronic system, each of which can have an arbitrary kinetic energy up to some maximum value. Therefore the complete kinematics of DIS is determined by three independent invariants rather than two as we are used to in elastic collisions. A natural choice of one of these invariants is the square of the total CMS energy  $S$ ,

$$S = (k + P)^2 = m_p^2 + 2k \cdot P \quad (86)$$

which is defined by the beam energy.

The second invariant is usually chosen to be the negative square of 4-momentum transfer:

$$Q^2 = -q^2 = -(k - k')^2 = 4EE' \sin^2 \frac{\theta}{2} \quad (87)$$

The third independent invariant can be taken to be  $W$  or alternatively one of the dimensionless variables

$$x = \frac{Q^2}{2P \cdot q} \quad (88)$$

or

$$y = \frac{P \cdot q}{k \cdot P} \quad (89)$$

where  $q = k - k'$ .

The variable  $y$  has a simple physical meaning in the target rest frame where  $P = (m_p, 0, 0, 0)$ ,  $k = (E_{\text{LAB}}, 0, 0, E_{\text{LAB}})$ , and  $k' = (E'_{\text{LAB}}, \vec{p}_3)$ , hence  $y = 1 - E'_{\text{LAB}}/E_{\text{LAB}}$ , *i.e.*  $y$  is the relative energy loss of the electron in the LAB frame.

The invariant  $x$  is the Bjorken scaling variable or simply Bjorken- $x$ . It was first recognised as an important variable of DIS by J.D. Bjorken who predicted the property of scaling in DIS which was subsequently confirmed experimentally.

Interesting is the expression of  $S$  in terms of the beam energies. In fixed target DIS we have the electron or muon beam with 4-momentum  $k = (E, 0, 0, E)$  and the proton target with  $P = (m_p, 0, 0, 0)$ , hence

$$S = m_p^2 + 2m_p E$$

whereas in an electron-proton collider like HERA we have 4-momenta  $P = (E_p, 0, 0, E_p)$  and  $k = (E_e, 0, 0, -E_e)$  and hence

$$S = 4E_e E_p$$

# Kinematics of DIS - 3

Other useful relations between the various kinematical variables are the following:

$$Q^2 = xyS \quad (90)$$

and

$$W^2 = m_p^2 + Q^2(1/x - 1) \quad (91)$$

where in the latter formula we have kept the proton mass in order to indicate that the threshold of  $W$  corresponds to elastic scattering.

Within the framework of the parton model, DIS proceeds by the exchange of a photon or intermediate vector boson with only one of the quarks in the proton. This is shown in the diagram in Fig. 3.

The electron-quark collision is elastic. As a result of this collision the struck quark acquires a sufficient momentum to break away from the rest of the proton as far as the colour force allows it to travel. At this stage some of the binding energy is converted into the creation of a quark-antiquark pair from the vacuum; the antiquark combines with the original quark into a meson, leaving behind a quark which can give rise to the creation of another quark-antiquark




Figure 3: Parton model diagram of deep inelastic scattering.



# Kinematics of DIS - 4

pair. This process, called *fragmentation*, continues until the remaining energy drops below the threshold for the creation of another pair. Thus, as a result of fragmentation, several mesons are created which travel roughly in the direction of the struck quark. Such a system of mesons, or more generally of hadrons, is called a *jet*. The residue of the proton is a highly unstable system: it has lost a quark, absorbed a quark presumably of the wrong sort that is left over from the fragmentation, and has absorbed a fraction of the energy transferred from the electron. Therefore it breaks up into several hadrons.

The elastic electron-quark collision is the *hard subprocess* of DIS. If we think of the incoming electron and proton as travelling in opposite directions, then the quark carries a fraction of the proton momentum. At a sufficiently high momentum, where the proton mass is negligible, the energy of the quark is the same fraction of the proton energy. It turns out that this fraction is identical with the Bjorken- $x$  defined above. Denoting the 4-momentum of the incoming quark by  $p$  we have therefore

$$p = xP$$

Denoting the invariant  $(k + p)^2$  by  $s$ , which is the squared CMS energy of the subprocess, we have therefore also

$$s = xS \tag{92}$$

This, together with the definition of  $Q^2$ , shows that the two independent invariants that control the kinematics of the subprocess are  $x$  and  $Q^2$ .



# Kinematics of DIS - 5

The first DIS experiments were carried out in 1967 at the Stanford 2-mile linear electron accelerator with electron beams of up to 20 GeV and hydrogen targets at rest, giving a CMS energy of about 6 GeV. Subsequent fixed target experiments were done in other laboratories, notably at the CERN SPS with muon beams of up to nearly 300 GeV and hence of CMS energies up to about 25 GeV. The range of energies available for DIS was extended by an order of magnitude when in 1992 the electron-proton collider HERA came into operation at the DESY laboratory in Hamburg. In this collider the electrons are accelerated up to nearly 30 GeV and the protons up to 820 GeV, giving a CMS energy of 314 GeV. Theoretically the corresponding values of  $Q^2$  go up to about  $10^5 \text{ GeV}^2$ .

An important tool to study the structure of the nucleon is also deep inelastic scattering with neutrinos as beam particles. The kinematics is identical with the one described above, but one must bare in mind that the exchanged particle in neutrino-DIS is an intermediate vector boson, either the  $W$  or the  $Z$ .





# $F_1$ and $F_2$

$$F_1(x, Q^2) \rightarrow \frac{1}{2} \sum_f Q_f^2 (q_f(x) + \bar{q}_f(x)). \quad (28)$$

The result in Eq. (28) demonstrates that  $F_1$  depends only on the dimensionless variable  $x = Q^2/2\nu$  in the deep inelastic limit, which is known as Bjorken scaling[5, 6]. The experimental observation of this scaling was the first direct evidence for point-like constituents in hadrons[7]. The quark distribution functions  $q_f(x)$ ,  $\bar{q}_f(x)$  defined by Eq. (26) for  $x \geq 0$  are an intrinsic non-perturbative property of the

hadron  $H$ . They may be interpreted as momentum distributions for quarks and anti-quarks inside the hadron and in principle (though not yet in practice) they can be computed from a non-perturbative analysis in QCD. At present these distribution functions must simply be determined experimentally from (largely) DIS experiments. We also find that

$$F_2(x, Q^2) = 2xF_1(x, Q^2) = x \sum_f Q_f^2 (q_f(x) + \bar{q}_f(x)). \quad (29)$$

The form of the relation between  $F_1$  and  $F_2$  is a consequence of the spin 1/2 nature of the struck quark. The difference is proportional to the longitudinal structure function  $F_L(x, Q^2)$ , and is zero at lowest order due to helicity conservation[8].

Applying these results to deep inelastic scattering on a proton target the proton wavefunction is dominated by  $uud + \dots$  where the dots indicate  $uud$  plus further quarks (including heavy flavours). With notation  $q_u(x) = u(x)$ ,  $\bar{q}_u(x) = \bar{u}(x)$  etc,

$$F_{2,\text{proton}}(x, Q^2) \sim x \left( \frac{4}{9}(u(x) + \bar{u}(x)) + \frac{1}{9}(d(x) + \bar{d}(x)) + \text{heavy flavours} \right). \quad (30)$$

We note that the derivation of Eq. (28) is an approximation which relies on the assumption that  $k$ , being the momentum of a quark (or antiquark) inside the proton, should have a very small probability of having any momentum components greater than  $\mathcal{O}(\Lambda_{QCD})$ . As such it also implies corrections of  $\mathcal{O}(\Lambda_{QCD}^2/Q^2)$  corresponding to higher twist operators (as discussed in[9]). However, it also ignores







# Hera

Reduced uncertainty

HERA I




HERA I + II






# Proton Parton Densities

H1 and ZEUS HERA I+II Combined PDF Fit



H1 and ZEUS Combined PDF Fit




HERA Structure Functions Working Group July 2010




# Proton Parton Densities

H1 and ZEUS HERA I+II Combined PDF Fit



H1 and ZEUS Combined PDF Fit




HERA Structure Functions Working Group July 2010



# Parton Distributions @ $Q^2 = 10 \text{ TeV GeV}$



NNPDF2.3 NNLO  $N_F = 6$





# Scaling violations in DIS

Increasing  $Q^2$  probes inner structure of proton  
 High  $x$ : quarks      Low  $x$ : gluons



At low  $x$  we have gluon  
 at high  $x$  we have quarks

Quarks: when  $Q^2$  increases  $F_2(x)$  goes down



Gluons: when  $Q^2$  increases  $F_2(x)$  Goes up






# Scaling Violations [SLAC 1972]

$$F_2(x, Q^2) = \sum e_q^2 x q(x, Q^2)$$





$F_2$   
[ $\times 2^i$ ]



## Scaling Violations

[HERA & fixed target data]

Precision: 2-3%  
[bulk region]

For  $x < 10^{-2}$ :

$$\frac{dF_2}{d\log Q^2} \sim g(x, Q^2) \cdot \alpha_s(Q^2)$$

NLO QCD Fits:


Quark densities  
Gluon density  
Strong coupling  $\alpha_s$



# Which region $x$ - $Q^2$ is seen by different experiments?

$$W^2 = m_p^2 + Q^2(1/x - 1)$$

mass of the recoiling quark  
against the lepton









# Particle Production @ LHC

## LHC parton kinematics



$pp \rightarrow X_M + \text{remnants}$

$X_M$ : particle with mass  $M$   
 e.g. Higgs

$$M^2 = x_1 x_2 \cdot s$$

i.e. to produce a particle with mass  $M$   
 at LHC energies ( $\sqrt{s} = 14 \text{ TeV}$ )

$$\langle x \rangle = \sqrt{x_1 x_2} = M/\sqrt{s}$$

[ $x_1 = x_2$ : mid-rapidity]


LHC needs:

Knowledge of parton densities  
 Extrapolation over orders of magnitudes





# Kinematic domains in DIS



- $x$  and  $Q^2$  domains probed by fixed-target and collider experiments.
- Some of the final states accessible at the LHC are indicated
- $Q^2 = M^2$  is the mass of some states accessible to LHC.
- For example, exclusive  $J/\psi$  and upsilon production at high  $|y|$  at the LHC may probe/imply the gluon PDF down to  $x \sim 10^{-5}$ .



# DGLAP Equations: extrapolating $F_2$

[ $z$ : momentum fraction of radiated parton]

[DGLAP: Dokshitzer, Gribov, Lipatov, Altarelli, Parisi]

$$\frac{\partial}{\partial \log Q^2} \begin{bmatrix} q(x, Q^2) \\ g(x, Q^2) \end{bmatrix} = \frac{\alpha_s}{2\pi} \begin{bmatrix} P_{q/q} \left[ \begin{array}{c} \gamma \\ x \end{array} \right] & P_{q/g} \left[ \begin{array}{c} \gamma \\ x \end{array} \right] \\ P_{g/q} \left[ \begin{array}{c} \gamma \\ x \end{array} \right] & P_{g/g} \left[ \begin{array}{c} \gamma \\ x \end{array} \right] \end{bmatrix} \otimes \begin{bmatrix} q(x, Q^2) \\ g(x, Q^2) \end{bmatrix}$$

PDFs

$$\frac{4}{3} \left[ \frac{1+z^2}{1-z} \right] \quad \frac{1}{2} [z^2 + (1-z^2)] \quad \frac{4}{3} \left[ \frac{1+(1-z)^2}{z} \right]$$

$$P \otimes f(x, Q^2) = \int_x^1 \frac{dy}{y} P(x/y) f(y, Q^2)$$

$$6 \left[ \frac{z}{1-z} + \frac{1-z}{z} + z(1-z) \right]$$



# Measurement of parton functions: which data?

**Table 19.1:** The main processes relevant to global PDF analyses, ordered in three groups: fixed-target experiments, HERA and the  $p\bar{p}$  Tevatron /  $pp$  LHC. For each process we give an indication of their dominant partonic subprocesses, the primary partons which are probed and the approximate range of  $x$  constrained by the data.

Fixed – target experiments

HERA & Tevatron

LHC



Process	Subprocess	Partons	$x$ range
$\ell^\pm \{p, n\} \rightarrow \ell^\pm X$	$\gamma^* q \rightarrow q$	$q, \bar{q}, g$	$x \gtrsim 0.01$
$\ell^\pm n/p \rightarrow \ell^\pm X$	$\gamma^* d/u \rightarrow d/u$	$d/u$	$x \gtrsim 0.01$
$pp \rightarrow \mu^+ \mu^- X$	$u\bar{u}, d\bar{d} \rightarrow \gamma^*$	$\bar{q}$	$0.015 \lesssim x \lesssim 0.35$
$pn/pp \rightarrow \mu^+ \mu^- X$	$(u\bar{d})/(u\bar{u}) \rightarrow \gamma^*$	$\bar{d}/\bar{u}$	$0.015 \lesssim x \lesssim 0.35$
$\nu(\bar{\nu}) N \rightarrow \mu^-(\mu^+) X$	$W^* q \rightarrow q'$	$q, \bar{q}$	$0.01 \lesssim x \lesssim 0.5$
$\nu N \rightarrow \mu^- \mu^+ X$	$W^* s \rightarrow c$	$s$	$0.01 \lesssim x \lesssim 0.2$
$\bar{\nu} N \rightarrow \mu^+ \mu^- X$	$W^* \bar{s} \rightarrow \bar{c}$	$\bar{s}$	$0.01 \lesssim x \lesssim 0.2$
$e^\pm p \rightarrow e^\pm X$	$\gamma^* q \rightarrow q$	$g, q, \bar{q}$	$10^{-4} \lesssim x \lesssim 0.1$
$e^+ p \rightarrow \bar{\nu} X$	$W^+ \{d, s\} \rightarrow \{u, c\}$	$d, s$	$x \gtrsim 0.01$
$e^\pm p \rightarrow e^\pm c\bar{c}X, e^\pm b\bar{b}X$	$\gamma^* c \rightarrow c, \gamma^* g \rightarrow c\bar{c}$	$c, b, g$	$10^{-4} \lesssim x \lesssim 0.01$
$e^\pm p \rightarrow \text{jet}+X$	$\gamma^* g \rightarrow q\bar{q}$	$g$	$0.01 \lesssim x \lesssim 0.1$
$p\bar{p}, pp \rightarrow \text{jet}+X$	$gg, qg, qq \rightarrow 2j$	$g, q$	$0.00005 \lesssim x \lesssim 0.5$
$p\bar{p} \rightarrow (W^\pm \rightarrow \ell^\pm \nu) X$	$ud \rightarrow W^+, \bar{u}\bar{d} \rightarrow W^-$	$u, d, \bar{u}, \bar{d}$	$x \gtrsim 0.05$
$pp \rightarrow (W^\pm \rightarrow \ell^\pm \nu) X$	$u\bar{d} \rightarrow W^+, d\bar{u} \rightarrow W^-$	$u, d, \bar{u}, \bar{d}, g$	$x \gtrsim 0.001$
$p\bar{p}(pp) \rightarrow (Z \rightarrow \ell^+ \ell^-) X$	$uu, dd, ..(u\bar{u}, ..) \rightarrow Z$	$u, d, ..(g)$	$x \gtrsim 0.001$
$pp \rightarrow W^- c, W^+ \bar{c}$	$gs \rightarrow W^- c$	$s, \bar{s}$	$x \sim 0.01$
$pp \rightarrow (\gamma^* \rightarrow \ell^+ \ell^-) X$	$u\bar{u}, d\bar{d}, .. \rightarrow \gamma^*$	$\bar{q}, g$	$x \gtrsim 10^{-5}$
$pp \rightarrow (\gamma^* \rightarrow \ell^+ \ell^-) X$	$u\gamma, d\gamma, .. \rightarrow \gamma^*$	$\gamma$	$x \gtrsim 10^{-2}$
$pp \rightarrow b\bar{b}X, t\bar{t}X$	$gg \rightarrow b\bar{b}, t\bar{t}$	$g$	$x \gtrsim 10^{-5}, 10^{-2}$
$pp \rightarrow \text{exclusive } J/\psi, \Upsilon$	$\gamma^*(gg) \rightarrow J/\psi, \Upsilon$	$g$	$x \gtrsim 10^{-5}, 10^{-4}$
$pp \rightarrow \gamma X$	$gq \rightarrow \gamma q, g\bar{q} \rightarrow \gamma\bar{q}$	$g$	$x \gtrsim 0.005$



# Backup



# Higgs Cross Section



The left column shows absolute results, the central column results normalized to the MSTW08 result, and the right column results normalized to each group's central result.

10 – 20 % PDF uncertainty

$m_H$




# DIS cross-section for Charged and Neutral Currents

$$d\sigma^2/dxdy$$

$$\eta_\gamma = 1; \quad \eta_{\gamma Z} = \left( \frac{G_F M_Z^2}{2\sqrt{2}\pi\alpha} \right) \left( \frac{Q^2}{Q^2 + M_Z^2} \right);$$

$$\eta_Z = \eta_{\gamma Z}^2; \quad \eta_W = \frac{1}{2} \left( \frac{G_F M_W^2}{4\pi\alpha} \frac{Q^2}{Q^2 + M_W^2} \right)^2.$$



The cross sections for neutral- and charged-current deep inelastic scattering on unpolarized nucleons can be written in terms of the structure functions in the generic form

$x$  = fraction of momentum carried by struck quark  
 $y$  = fraction of lepton energy lost

$$\frac{d^2\sigma^i}{dxdy} = \frac{4\pi\alpha^2}{xyQ^2} \eta^i \left\{ \left( 1 - y - \frac{x^2 y^2 M^2}{Q^2} \right) F_2^i + y^2 x F_1^i \mp \left( y - \frac{y^2}{2} \right) x F_3^i \right\}, \quad (19.8)$$

where  $i = \text{NC}$ , CC corresponds to neutral-current ( $eN \rightarrow eX$ ) or charged-current ( $eN \rightarrow \nu X$  or  $\nu N \rightarrow eX$ ) processes, respectively.

@ HERA